

Instruction for use

Intibio PRP (2ml)

Product composition

2ml of equine leukocyte and erythrocyte poor plasma containing physiological levels of thrombocytes.

Intended Use

Intibio PRP is an animal byproduct intended for use only by those who have received adequate training and knowledge on how to use orthobiologics in horses. A veterinary clinical follow-up is recommended 4 to 6 weeks after application.

Product Appearance

PRP is presented in a 2ml tube with container closure. The blue cap needs to be removed before aspiration of the product through the pierceable membrane.

How is Intibio PRP Prepared?

Each batch is tested for purity (thrombocytes, total protein, albumin), impurities (white blood cells, red blood cells), sterility (aerobic bacteria, anaerobic bacteria, fungi and yeast), conformity (colour, label inspection) and infectious diseases (over 20 different pathogens).

Storage Instructions

Intibio PRP is delivered as a 2ml tube and can be stored at:

- ◆ 4°C for 4 weeks
- ◆ -20°C (normal freezer) OR -80°C (specialist freezer) OR in liquid nitrogen (in the gas phase) for 24 months after production date

Intibio PRP can be refrozen once within 24 hours of defrosting. Freeze in an upright position to avoid liquid in the cap of the tube.

Due to the conditions of shipping, Intibio PRP may arrive with you partially defrosted. In this instance you can refreeze within 24hrs as above OR if you anticipate using the product within 4 weeks the tube can be stored at 4°C in the fridge.

Handling Instructions

When stored frozen, the product needs to be thawed in the palm of your hand or a warm water bath and heated to 37°C (body temperature) before use. Blood protein clusters or plasma crystals need to dissolve after stirring with a standard vortex.

When stored at 4°C, the product needs to be heated to body temperature before application. Blood protein clusters need to dissolve after stirring with a standard vortex.

Please make sure the tube cap remains closed at all times and check the content for a clear, yellow 2ml solution per tube before and after aspiration. In case where precipitations are noticed, make sure it is properly heated (37°C) and swirl the tube before aspiration.

Instruction for use

Intibio PRP (2ml)

Administration Instructions

Use sterile gloves and aseptically prepare the application site prior to use.

Disposal Instructions

Intibio PRP needs to be disposed as medical waste.

Additional Information/warnings

- ◆ When the content is not transparent or yellow or when the tube or cap is damaged, please do not apply the product and contact the operator.
More advanced pathology may result in an increased response after use.
- ◆ Do not mix with other products from animal origin or from different donors.
- ◆ When biological products from different donors are used in the same region a wash-out period of 4 weeks needs to be respected.
- ◆ Do not add more volume or other products in the tube as too much pressure may result in product leakage.
- ◆ Intended for veterinary use only.

